

Many ω -categorical Structures Have the Small Index Property

BERNHARD HERWIG

Paris, France

June 1994

Abstract

Theorem: Let A be a finite $\mathfrak{K}_m$ -free graph, $p_1, \dots, p_n$ partial isomorphisms of A . Then there exists a finite extension B , which is also a $\mathfrak{K}_m$ -free graph, and automorphisms f_i of B extending the p_i 's. This theorem can be used to prove the small index property for the generic countable graph of this class. The same method also works for a certain class of continuum many non isomorphic ω -categorical countable digraphs.

Hrushovski proved this theorem for the class of all finite graphs [Hr]; the proof presented here is an extension of his proof.

Notation: Let p be a partial mapping on a set A . By $D(p)$ we denote the domain of p by $R(p)$ the range of p (so $D(p) \subset A$ and $R(p) \subset A$). In this paper the partial mappings under consideration will always be injective. The edge relation will always be called R ; in the first part of the paper we will deal with graphs, so R will be symmetric and irreflexive, in the second part we will handle digraphs, so R will be antisymmetric and irreflexive.

Definition: If A is a graph and a is a point (possibly element of a graph extending A) we denote by $N_A(a)$ the set of neighbours of a , sometimes considered as a pure set and sometimes considered as a subgraph of A : $N_A(a) := \{b \in A \mid aRb\}$. A graph A is called $\mathfrak{K}_m$ -free (for $m \in \omega$), if $\mathfrak{K}_m$, which is the complete graph with m vertices, is not embeddable into A , i.e. there does not exist $a_1, \dots, a_m \in A$ such that a_iRa_j (for $1 \leq k < l \leq m$).

Theorem 1 *Let $m, n \geq 1$, let A be a finite $\mathfrak{K}_m$ -free graph. Let $p_1, \dots, p_n$ be partial isomorphisms on A . There exists a finite $\mathfrak{K}_m$ -free graph B , $B \supset A$, and $f_1, \dots, f_n \in \text{Aut}(B)$, such that $f_i \supset p_i$.*

Before going through the formal proof, which looks a little bit technical, let us describe the main ideas of the proof:

We know that the theorem holds for $\mathfrak{K}_3$ -free graphs [Hg]. The proof in the general case will be by induction on m . Let us introduce for every $a \in A$ a new colour (i.e. a unary predicate) U_a such that for $b \in A$ b is of colour U_a iff bRa . Now A is $\mathfrak{K}_m$ -free if and only if you can not embed $\mathfrak{K}_{m-1}$ “uni-coloured” into A , that means there does not exist a colour U_a and elements $a_1, \dots, a_{m-1}$ in A such that $a_k Ra_l$ ($1 \leq k < l \leq m$) and all the a_i ’s are of colour U_a . Thus one can reduce $\mathfrak{K}_m$ -freeness conditions to certain $\mathfrak{K}_{m-1}$ -freeness conditions, if one works with coloured graphs. Here are the main problems, one has to overcome doing this reduction.

1. With respect to the colours p_i is not longer a partial isomorphism, it is a “permorphism”, i.e. it respects the colours only up to a permutation χ_i of the colours ($(U_a)^{\chi_i} = U_{a^{p_i}}$). But as can be seen in [Hg] Hrushovski’s original proof of the theorem in the case of the graphs [Hr] also works for permorphisms.
2. χ_i is not yet really a permutation of the set of colours $\{U_a \mid a \in A\}$, it is only a partial function. As in [Hg] one overcomes this problem by doing a type realizing step to get a nice graph $C \supset A$, afterwards one looks at the colours $\{U_d \mid d \in C\}$ and extends the partially already defined functions χ_i to permutations of this set.
3. If one extends the graph A considered as a (uni-coloured $\mathfrak{K}_{m-1}$)-free graph to a (uni-coloured $\mathfrak{K}_{m-1}$)-free graph B and the p_i ’s to f_i ’s, how can one ensure that B is $\mathfrak{K}_m$ -free? Take into account that B is $\{U_d \mid d \in C\}$ -coloured, so we don’t have for every $b \in B$ a colour U_b such that all neighbours of b have colour U_b . This problem disappears by miracle: The resulting graph B looks locally like A . Especially for every $a \in A$ the neighbours of a in B will still all have colour U_a . Now any orbit of the automorphism group of B (as graph) will have an element inside A . So to check $\mathfrak{K}_m$ -freeness of B , one has only to look for copies of $\mathfrak{K}_m$ having one element a inside A , but such a copy would lead to a copy of $\mathfrak{K}_{m-1}$ of colour U_a .
4. To get a proper induction, one has to prove the theorem for coloured graphs and permorphisms. Starting with a coloured graph, we have to introduce a new set of colours $\{U_d \mid d \in C\}$. But (uni-coloured $\mathfrak{K}_m$)-freeness does not exactly mean (uni-coloured $\mathfrak{K}_{m-1}$)-freeness with

respect to the new colours. It means precisely: there does not exist an old colour U and a new colour U_a such that a is of colour U and a U_a -coloured copy of $\mathfrak{K}_{m-1}$ which is at the same time U -coloured. We will call such a combination (U, U_a) a critical colouring, and we have to avoid copies of $\mathfrak{K}_{m-1}$ which are critical coloured. This last problem and the notational complication arising from the fact that we are dealing with permorphisms rather than isomorphisms will make the prove look rather technical.

The graph $C \supset A$ we will get by the type realizing step will have as additional feature that every element in A has exactly the same number of neighbours in C , i.e. the same number of colours. In the proof we will maintain this condition, even if this is not really necessary. If one erases in the proof all statements saying something like “ $\#\mathcal{U}^j(a) = d_j$ ” one get a slightly shorter proof.

Now the definitions which follow, and the version of the theorem (i.e. Lemma 2) which will be provable by induction should be sufficiently motivated.

Definition: Let S be a relational language. Let χ be a permutation of the symbols in S mapping every symbol to a symbol with the same arity. Let A be a S -structure and p be a partial mapping on A . p is called a χ -permorphism, if for every $r \in \omega$ and every r -ary relation R in S and every $a_1, \dots, a_r \in D(p)$: $Ra_1 \dots a_r \iff R^\chi a_1^p \dots a_r^p$.

Definition: Let $\mathcal{U}^1, \dots, \mathcal{U}^r$ be a family of disjoint finite sets of unary predicates (called colours) and $d_1, \dots, d_r$ be constants. Let $\mathcal{U} := \bigcup_{1 \leq j \leq r} \mathcal{U}^j$, $\mathcal{L} := \mathcal{U} \cup \{R\}$. If A is a $\mathcal{L}$ -structure and $a \in A$, $V \in \mathcal{U}$, then we write $a \in V$ to indicate that the unary predicate V (or rather its interpretation in A) is true for a ; we also write “ a is of colour V ”.

1. A $\mathcal{U}$ -graph is a $\{R\} \cup \mathcal{U}$ -structure A such that A considered as a $\{R\}$ -structure is a graph and furthermore for every $a \in A$ and every j ($1 \leq j \leq r$): $\#\mathcal{U}^j(a) = d_j$. Here $\mathcal{U}^j(a) := \{V \in \mathcal{U}^j \mid a \in V\}$ is the set of colours of a . $\mathcal{U}(a) := \{V \in \mathcal{U} \mid a \in V\}$.
2. Let $\mathcal{U}_c \subset \mathcal{U}^1 \times \dots \times \mathcal{U}^r$. $\mathcal{U}_c$ will be called the set of critical colourings. We call A $\mathcal{U}_c$ - $\mathfrak{K}_m$ -free (for $m \in \omega$), if there does not exist a colouring

$(V_1, \dots, V_r) \in \mathbf{U}_c$ and elements $a_1, \dots, a_m \in A$, such that $a_k R a_l$ (for $1 \leq k < l \leq m$) and $a_k \in V_j$ (for $1 \leq k \leq m$, $1 \leq j \leq r$).

Lemma 2 *Let $m \geq 1$. Let $\mathbf{U}^1, \dots, \mathbf{U}^r$ be disjoint sets of colours (where $r \geq 0$). Let $\mathbf{U} := \bigcup_{1 \leq j \leq r} \mathbf{U}^j$. Let $\chi_i^j \in \text{Sym}(\mathbf{U}^j)$ (for $1 \leq i \leq n$, $1 \leq j \leq r$), $\mathcal{L} := \mathbf{U} \cup \{R\}$ $\chi_i := \bigcup_{0 \leq j \leq r} \chi_i^j \in \text{Sym}(\mathcal{L})$, where χ_i^0 is the identity on $\{R\}$. Let furthermore $\mathbf{U}_c \subset \mathbf{U}^1 \times \dots \times \mathbf{U}^r$ be a set called critical colourings. Suppose $\mathbf{U}_c$ is χ_i -invariant (for $1 \leq i \leq n$).*

Let A be a $\mathbf{U}$ -graph. We suppose that for every $1 \leq j \leq r$ there is given a constant d_j , such that for every $a \in A$ $\#(\mathbf{U}^j(a)) = d_j$. Suppose A is $\mathbf{U}_c$ - $\mathcal{K}_m$ -free.

Let $p_1, \dots, p_n$ be partial mappings on A such that p_i is a χ_i -permorphism. We suppose further that for $a \in A$ and $1 \leq j \leq r$ there exists a colour $U_a^j \in \mathbf{U}^j$ such that for $b \in A$: $b \in U_a^j \iff a R b$ and such that for $a \in D_i = D(p_i)$: $(U_a^j)^{\chi_i} = U_{a^{p_i}}^j$. Let us finally suppose that for every $a, b \in A$, if $a \neq b$ then there exist an i ($1 \leq i \leq n$) such that $a^{p_i} = b$. (This is not really necessary.)

Then there exist a $\mathbf{U}$ -graph B , $B \supset A$, B $\mathbf{U}_c$ - $\mathcal{K}_m$ -free, $f_1, \dots, f_n \in \text{Aut}(B)$, $f_i \supset p_i$, f_i a χ_i -permorphism (for $1 \leq i \leq n$). B satisfies in addition: $\forall b \in B \forall a \in A : a R b \Rightarrow \forall j (1 \leq j \leq r) b \in U_a^j$. Furthermore B will be chosen to satisfy in addition: $\forall b \in B \exists f \in \langle f_1, \dots, f_n \rangle : b^f \in A$.

From the Lemma follows the theorem:

Let in the Lemma $r = 0$, that means we just talk about (uncoloured) graphs. $\mathbf{U}_1 \times \dots \times \mathbf{U}_r$ just contains the empty tuple λ and we let $\mathbf{U}_c = \{\lambda\}$, then $\mathbf{U}_c$ - $\mathcal{K}_n$ -freeness just means $\mathcal{K}_n$ -freeness. χ_i is the identity on $\{R\}$ and χ_i -permorphism just means isomorphism of graphs. We can suppose w.l.o.g. that for every $a, b \in A$ there exists an i ($1 \leq i \leq n$) such that $a^{p_i} = b$ for example by assuming that $\{p_1, \dots, p_n\}$ equals the set of all partial isomorphisms on A . Thus the lemma for $r = 0$ just implies the theorem.

Proof of the Lemma:

The proof goes by induction on m . Lets first treat the case $m = 1$.

The proof follows in this case the lines of the original proof of Hrushovski

[Hr]. The first step will be a type realizing step. A subset of A determines a “type over A ” (considered as a pure graph) in our context.

Claim:

- a) There exist a finite graph C , $C \supset A$ (C is a pure graph so $C \supset A$ means A is a substructure of C as graphs) and a constant c_0 such that for every $A_0 \subset A$: $\#\{c \in C \mid N_A(c) = A_0\} = c_0$.
- b) There exist bijections $h_1, \dots, h_n \in \text{Sym}(C)$, $h_i \supset p_i$, such that for every $a \in D_i$, $b \in C$: $aRb \iff a^{p_i}Rb^{h_i}$.

Proof of a):

Let for $A_0 \subset A$ $c_{A_0} := \#\{c \in A \mid N_A(c) = A_0\}$ and $c_0 := \max\{c_{A_0} \mid A_0 \subset A\}$. Now for every $A_0 \subset A$ add $(c_0 - c_{A_0})$ many points which have as set of neighbours A_0 to get C .

Proof of b):

Fix i ($1 \leq i \leq n$) We first check, that for every $D_0 \subset D_i$ ($D_i := D(p_i)$, $R_i := R(p_i)$)

$$\#(\{c \in C \mid N_{D_i}(c) = D_0\}) = \#(\{c \in C \mid N_{R_i}(c) = D_0^{p_i}\}):$$

$$\#(\{c \in C \mid N_{D_i}(c) = D_0\}) = \sum_{B \subset (A - D_i)} \#(\{c \in C \mid N_A(c) = D_0 \cup B\}) =$$

$$c_0 \cdot 2^{\#(A - D_i)} = c_0 \cdot 2^{\#(A - R_i)} = \#(\{c \in C \mid N_{R_i}(c) = D_0^{p_i}\}).$$

Furthermore, because p_i is a partial isomorphism of graphs, p_i maps $\{c \in C \mid N_{D_i}(c) = D_0\} \cap D_i$ to $\{c \in C \mid N_{R_i}(c) = D_0^{p_i}\} \cap R_i$. So we find $h_i \supset p_i$ a bijection of C mapping for every $D_0 \subset D_i$ $\{c \in C \mid N_{D_i}(c) = D_0\}$ to $\{c \in C \mid N_{R_i}(c) = D_0^{p_i}\}$, but this means exactly that h_i has the property we want. ■(Claim)

Now we do a duplicator step:

We fix C and $h_1, \dots, h_n$, which we get from the claim. We let $\Gamma \subset \text{Sym}(\mathcal{L}) \times \text{Sym}(C)$ be the subgroup generated by the elements $\gamma_i = (\chi_i, h_i)$ ($1 \leq i \leq n$). Note that if $\gamma = (\chi, h) \in \Gamma$ then χ fixes $\mathbf{U}^j$ (for $1 \leq j \leq r$) and χ fixes $\mathbf{U}_c$. On $A \times \Gamma$, we define the equivalence relation $\equiv$ to be the symmetric, reflexive and transitive closure of $E = \{((a^{p_i}, \gamma), (a, \gamma_i \gamma)) \mid 1 \leq i \leq n, a \in D_i, \gamma \in \Gamma\}$.

We note some basic facts:

1. If $(a, (\chi, h)) \equiv (a_s, (\chi_s, h_s))$ then $a^h = a_s^{h_s}$ and $(\mathbf{U}(a))^x = (\mathbf{U}(a_s))^{\chi_s}$

2. If $(a, (\chi, h)) \equiv (a_s, (\chi_s, h_s))$ and $(b, (\chi, h)) \equiv (b_s, (\chi_s, h_s))$ then
 $aRb \iff a_sRb_s$
3. If $(a, (\chi, h)) \equiv (a_s, (\chi_s, h_s))$ and $c \in C$ then
 $aRc^{h^{-1}} \iff a_sRc^{h_s^{-1}}$ and $(U_a^j)^\chi = (U_{a_s}^j)^{\chi_s}$.

Proof of the facts:

1. It suffices to prove 1. in the case $((a, (\chi, h)), (a_s, (\chi_s, h_s)))$ is actually in E , so $a_s \in D_i$ and $a_s^{p_i} = a$ and $(\chi_s, h_s) = (\chi_i, h_i) \cdot (\chi, h)$. But then $a^h = a_s^{p_i h} = a_s^{h_i h} = a_s^{h_s}$ and $\mathbf{U}(a)^\chi = \mathbf{U}(a_s^{p_i})^\chi = \mathbf{U}(a_s)^{\chi_i \chi} = \mathbf{U}(a_s)^{\chi_s}$ because p_i is a χ_i -permorphism.
3. Again we can suppose that $a_s \in D_i$, $a = a_s^{p_i}$ and $(\chi_s, h_s) = (\chi_i, h_i)(\chi, h)$. So $a_sRc^{h_s^{-1}} \iff a_sRc^{h^{-1}h_i^{-1}} \iff a_s^{p_i}Rc^{h^{-1}}$ by the condition (in claim b)) on h_i and $(U_{a_s}^j)^{\chi_s} = (U_{a_s}^j)^{\chi_i \chi} = (U_{a_s^{p_i}}^j)^\chi = (U_a^j)^\chi$ by the conditions on U_a^j in the hypothesis of the lemma.
2. $aRb \iff aRb^{hh^{-1}} \iff a_sRb^{hh_s^{-1}} \iff a_sRb_s^{h_s h_s^{-1}} \iff a_sRb_s$,
by 1. and 3.

Now we are ready to define a $\mathcal{L}$ -structure on $A \times \Gamma / \equiv$:

For $e, f \in A \times \Gamma / \equiv$:

$eRf \iff \exists \gamma \in \Gamma \exists a, b \in A \ e = (a, \gamma) / \equiv, \ f = (b, \gamma) / \equiv$ and aRb .

For $V \in \mathbf{U}$ and $e \in A \times \Gamma / \equiv$ we define

$e \in V \iff \exists (\chi, h) \in \Gamma \exists a \in A \ e = (a, (\chi, h)) / \equiv$ and $a \in V^{\chi^{-1}}$.

We note that

4. For $\gamma \in \Gamma \ a, b \in A$: $((a, \gamma) / \equiv) R ((b, \gamma) / \equiv) \iff aRb$.
5. For $(\chi, h) \in \Gamma, a \in A, V \in \mathbf{U}$: $((a, (\chi, h)) / \equiv) \in V^\chi \iff a \in V$

Proofs:

4. follows directly from 2.

5. follows from 1.:

$$(a, (\chi, h)) / \equiv \in V^\chi \iff$$

$$(\exists a_s, \chi_s, h_s \ (a, (\chi, h)) \equiv (a_s, (\chi_s, h_s)) \text{ and } a_s \in V^{\chi \chi_s^{-1}}) \iff a \in V,$$

$$\text{namely: } a_s \in V^{\chi \chi_s^{-1}} \iff V^{\chi \chi_s^{-1}} \in \mathbf{U}(a_s) \iff V^\chi \in \mathbf{U}(a_s)^{\chi_s} \iff$$

$$V^\chi \in \mathbf{U}(a)^\chi \iff V \in \mathbf{U}(a) \iff a \in V.$$

■(facts)

We define a map $i : A \rightarrow A \times \Gamma / \equiv$ by $i(a) := (a, 1) / \equiv$ (where 1 is the unit element in the group Γ). i is injective: if $(a, 1) \equiv (b, 1)$ then by 1. $a = b$. By 4. and 5. i is an embedding of A into $A \times \Gamma / \equiv$ as $\mathcal{L}$ -structures. By identifying we suppose $A \subset A \times \Gamma / \equiv$ and we set $B = A \times \Gamma / \equiv$. B is $\mathcal{U}_c\text{-}\mathcal{K}_1$ -free. For suppose there exists $(a, (\chi, h)) / \equiv \in B$ and $(V_1, \dots, V_r) \in \mathcal{U}_c$ such that $(a, (\chi, h)) / \equiv \in V_j$ (for $1 \leq j \leq r$), then by 5. $a \in V_j^{\chi^{-1}}$ and $(V_1^{\chi^{-1}}, \dots, V_r^{\chi^{-1}}) \in \mathcal{U}_c$, because χ^{-1} fixes $\mathcal{U}_c$. But this contradicts $\mathcal{U}_c\text{-}\mathcal{K}_1$ -freeness of A .

We define the automorphism f_i (for $1 \leq i \leq n$) by $((a, \gamma) / \equiv)^{f_i} := (a, \gamma \gamma_i) / \equiv$. This is an χ_i -permorphism, because for $V \in \mathcal{U}$ and $(a, (\chi, h)) / \equiv \in B$:

$$\begin{aligned} (a, (\chi, h)) / \equiv \in V &\iff a \in V^{\chi^{-1}} \iff a \in (V^{\chi_i})^{\chi_i^{-1} \chi^{-1}} \iff \\ &(a, (\chi \chi_i, h h_i)) / \equiv \in V^{\chi_i} \iff ((a, (\chi, h)) / \equiv)^{f_i} \in V^{\chi_i} \end{aligned}$$

The mapping $\varphi : \Gamma \rightarrow \langle f_1, \dots, f_n \rangle$ (sending γ_i to f_i), which is given by $((a, \gamma') / \equiv)^{\varphi(\gamma)} = (a, \gamma' \gamma) / \equiv$ is a surjective homomorphism of groups, and for $((a, \gamma) / \equiv) \in B$ we have $((a, \gamma) / \equiv)^{\varphi(\gamma^{-1})} = (a, 1) / \equiv \in A$.

Finally let $a \in A$, $b \in B$, $a R b$. So let $a = (a, 1) / \equiv = (a_s, (\chi_s, h_s)) / \equiv$ and $b = (b_s, (\chi_s, h_s)) / \equiv$. Because $a_s R b_s$ we have $b_s \in \mathcal{U}_{a_s}^j$ (in A). So by 5. $b \in (\mathcal{U}_{a_s}^j)^{\chi_s}$ but by 3. $(\mathcal{U}_{a_s}^j)^{\chi_s} = \mathcal{U}_{a_s}^j$. This finishes the case $m = 1$. $\blacksquare(m = 1)$

Now we do the step of induction $m \rightarrow m + 1$ ($m \geq 1$):

We have the set of colours $\mathcal{U}^1, \dots, \mathcal{U}^r$ and we know that A is $\mathcal{U}_c\text{-}\mathcal{K}_{m+1}$ -free. By a type realizing step and by introducing new colours we want to consider A as satisfying a certain $\mathcal{K}_m$ -freeness condition and then we want to apply the Lemma for m .

A subset $A_0 \subset A$ and a colouring $\mathcal{U}_0 \subset \mathcal{U}$ determines a type over A in this context. But not all of the types are realizable in $\mathcal{U}_c\text{-}\mathcal{K}_{m+1}$ -free graphs. We call a tuple $(A_0, \mathcal{U}_0)$ realisable, if for all j ($1 \leq j \leq r$) $\#(\mathcal{U}_0 \cap \mathcal{U}^j) = d_j$ and if there does not exist $(V_1, \dots, V_r) \in \mathcal{U}_c \cap \mathcal{U}_0 \times \dots \times \mathcal{U}_0$ and elements $a_1, \dots, a_m \in A_0$ such that $a_k R a_l$ (for $1 \leq k < l \leq m$) and $a_k \in V_j$ (for $1 \leq k \leq m$, $1 \leq j \leq r$). Here are facts about realizability:

1. If $B_0 \subset A_0 \subset A$ and if $(A_0, \mathcal{U}_0)$ is realisable, then $(B_0, \mathcal{U}_0)$ is realisable.
2. If $C \supset A$ (C a $\mathcal{U}$ -graph) is $\mathcal{U}_c\text{-}\mathcal{K}_{m+1}$ -free, then for every $c \in C$ $(N_A(c), \mathcal{U}(c))$ is realisable. In particular for every $a \in A$ $(N_A(a), \mathcal{U}(a))$ is realisable.

3. If $(1 \leq i \leq n)$ and $D_0 \subset D_i$ and $\mathbf{u}_0 \subset \mathbf{u}$: $(D_0, \mathbf{u}_0)$ is realisable iff $(D_0^{p_i}, \mathbf{u}_0^{x_i})$ is realisable.

Proof of 3.:

Suppose $(D_0, \mathbf{u}_0)$ is realisable. Now we have for all j , $(1 \leq j \leq r)$: $\#(\mathbf{u}_0^{x_i} \cap \mathbf{u}^j) = \#(\mathbf{u}_0 \cap \mathbf{u}^j)^{x_i} = d_j$. Suppose there exists $(V_1^{x_i}, \dots, V_r^{x_i}) \in \mathbf{u}_c \cap \mathbf{u}_0^{x_i} \times \dots \times \mathbf{u}_0^{x_i}$ and $a_1^{p_i}, \dots, a_m^{p_i} \in D_0^{p_i}$ such that $a_k^{p_i} R a_l^{p_i}$ (for $1 \leq k < l \leq m$) and $a_k^{p_i} \in V_j^{x_i}$ (for $1 \leq k \leq m$ and $1 \leq j \leq r$). But then $(V_1, \dots, V_r) \in \mathbf{u}_c \cap \mathbf{u}_0 \times \dots \times \mathbf{u}_0$ and $a_1, \dots, a_m \in D_0$ and $a_k R a_l$ and $a_k \in V_j$, because $\mathbf{u}_c$ is χ_i -invariant and because p_i is a χ_i -permorphism. This is in contradiction to the realizability of $(D_0, \mathbf{u}_0)$, therefore $(D_0^{p_i}, \mathbf{u}_0^{x_i})$ must be realisable. ■(facts)

Now we do the type realizing step:

Claim:

- a) There exists a $\mathbf{u}_c$ - $\mathcal{K}_{m+1}$ -free $\mathbf{u}$ -graph $C \supset A$ and for every t ($0 \leq t \leq \#A$) a constant c_t such that for every $A_0 \subset A$ and every $\mathbf{u}_0 \subset \mathbf{u}$:

$$\#\{c \in C \mid N_A(a) \supset A_0, \mathbf{u}(a) = \mathbf{u}_0\} = \begin{cases} c_{\#A_0} & \text{if } (A_0, \mathbf{u}_0) \text{ is realisable} \\ 0 & \text{otherwise} \end{cases}$$

- b) There exist bijections $h_1, \dots, h_n \in \text{Sym}(C)$, $h_i \supset p_i$ such that for every $V \in \mathbf{u}$, for every $b \in C$, for every i ($1 \leq i \leq n$): $b \in V \iff b^{h_i} \in V^{x_i}$ and such that for every $a \in D_i$, $b \in C$ $a R b \iff a^{p_i} R b^{h_i}$.

Proof of a):

Let $T = \#A$. We construct graphs $A = C_T \subset C_{T-1} \subset \dots \subset C_0 = C$ and constants $c_T, \dots, c_0$ such that for every t ($T \geq t \geq 0$) and for every $A_0 \subset A$ with $\#A_0 \geq t$ and for $\mathbf{u}_0 \subset \mathbf{u}$: $\#\{c \in C_t \mid N_A(a) \supset A_0, \mathbf{u}(a) = \mathbf{u}_0\} = c_{\#A_0}$ if $(A_0, \mathbf{u}_0)$ is realisable and such that C_t is $\mathbf{u}_c$ - $\mathcal{K}_{m+1}$ -free. We set $c_T = 0$. If c_r, C_r are already constructed (for $T \geq r \geq t$) and $t \geq 1$ then we will construct C_{t-1} by adding points which have exactly $t-1$ neighbours, all of them in A : For $A_0 \subset A$ with $\#A_0 = t-1$, and for $\mathbf{u}_0 \subset \mathbf{u}$ such that $(A_0, \mathbf{u}_0)$ is realisable we define $c_{A_0, \mathbf{u}_0} := \#\{c \in C_t \mid N_A(c) \supset A_0, \mathbf{u}(c) = \mathbf{u}_0\}$ and we define c_{t-1} to be the maximum of all these $c_{A_0, \mathbf{u}_0}$. Now to get C_{t-1} we add for every realisable $(A_0, \mathbf{u}_0)$ (with $\#A_0 = t-1$) $c_{t-1} - c_{A_0, \mathbf{u}_0}$ many points, which have as set of neighbours exactly A_0 and as set of colours $\mathbf{u}_0$. C_{t-1} is a $\mathbf{u}$ -graph and $\mathbf{u}_c$ - $\mathcal{K}_{m+1}$ -free and for every $A_0 \subset A$ (with $t-1 \leq \#A_0$), for every $\mathbf{u}_0 \subset \mathbf{u}$: $\#\{c \in C_{t-1} \mid N_A(c) \supset A_0, \mathbf{u}(c) = \mathbf{u}_0\} = c_{\#A_0}$ if $(A_0, \mathbf{u}_0)$ is realisable. This is

true, because for $A_0 \geq t$ we did not change the set in question (by going from C_t to C_{t-1}) and if $\#A_0 = t - 1$ then $\#\{c \in C_{t-1} \mid N_A(c) \supset A_0, \mathbf{u}(c) = \mathbf{u}_0\} = \#\{c \in C_t \mid N_A(c) \supset A_0, \mathbf{u}(c) = \mathbf{u}_0\} + (c_{t-1} - c_{A_0, \mathbf{u}_0}) = c_{A_0, \mathbf{u}_0} + (c_{t-1} - c_{A_0, \mathbf{u}_0}) = c_{t-1}$.

The proof of b) is similar as the proof of claim b) in the case $m = 1$. Here it is crucial to check, that for every $D_0 \subset D_i = D(p_i)$ and every $\mathbf{u}_0 \subset \mathbf{u}(R_i := R(p_i))$:

$$\#\{c \in C \mid N_{D_i}(c) = D_0, \mathbf{u}(c) = \mathbf{u}_0\} = \#\{c \in C \mid N_{R_i}(c) = D_0^{p_i}, \mathbf{u}(c) = \mathbf{u}_0^{x_i}\}$$

This is done by downwards induction on the size of D_0 . The step of induction is in the case $(D_0, \mathbf{u}_0)$ is realisable (otherwise both sets are empty; here we are using fact 3.: $(D_0, \mathbf{u}_0)$ is realisable $\iff (D_0^{p_i}, \mathbf{u}_0^{x_i})$ is realisable) as follows:

$$\begin{aligned} & \#\{c \in C \mid N_{D_i}(c) = D_0, \mathbf{u}(c) = \mathbf{u}_0\} \\ = & \#\{c \in C \mid N_A(c) \supset D_0, \mathbf{u}(c) = \mathbf{u}_0\} \\ & - \sum_{D_0 \subsetneq E \subset D_i} \#\{c \in C \mid N_{D_i}(c) = E, \mathbf{u}(c) = \mathbf{u}_0\} \\ = & c_{\#D_0} - \sum_{D_0 \subsetneq E \subset D_i} \#\{c \in C \mid N_{R_i}(c) = E^{p_i}, \mathbf{u}(c) = \mathbf{u}_0^{x_i}\} \\ = & \#\{c \in C \mid N_A(c) \supset D_0^{p_i}, \mathbf{u}(c) = \mathbf{u}_0^{x_i}\} \\ & - \sum_{D_0^{p_i} \subsetneq E' \subset D_i} \#\{c \in C \mid N_{R_i}(c) = E', \mathbf{u}(c) = \mathbf{u}_0^{x_i}\} \\ = & \#\{c \in C \mid N_{R_i}(c) = D_0^{p_i}, \mathbf{u}(c) = \mathbf{u}_0^{x_i}\} \end{aligned}$$

■(claim)

Now we introduce a new set of colours: $\mathbf{u}^{r+1} = \{U_d^{r+1} \mid d \in C\}$, where U_d^{r+1} is a new unary predicate for every $d \in C$. We define $\chi_i^{r+1} \in \text{Sym}(\mathbf{u}^{r+1})$ (for $1 \leq i \leq n$) by $(U_d^{r+1})^{\chi_i^{r+1}} := U_{d^{h_i}}^{r+1}$. We let $\mathbf{u}' := \mathbf{u} \cup \mathbf{u}^{r+1}$ and $\chi'_i := \chi_i \cup \chi_i^{r+1} \in \text{Sym}(\mathbf{u}' \cup \{R\})$. $\mathbf{u}'_c \subset \mathbf{u}'^1 \times \dots \times \mathbf{u}'^{r+1}$ is defined by $(V_1, \dots, V_r, U_d^{r+1}) \in \mathbf{u}'_c \iff (V_1, \dots, V_r) \in \mathbf{u}_c$ and $d \in V_j$ ($1 \leq j \leq r$) (in C). $\mathbf{u}'_c$ is χ'_i -invariant, because $\mathbf{u}_c$ is χ_i -invariant, because of the definition of χ_i^{r+1} and because of the property of h_i in claim b) ($b \in V \iff b^{h_i} \in V^{x_i}$).

The colours in $\mathbf{u}^{r+1}$ are in a natural way interpreted in A : for $a \in A$ and $d \in C$ we define $a \in U_d^{r+1} \iff dRa$ (in C). Now A is a $\mathbf{u}'$ -graph. We have

for every $a \in A$:

$$\begin{aligned} \#(\mathbf{u}^{r+1}(a)) &= \#(\{d \in C \mid a \in \mathbf{u}_d^{r+1}\}) = \#(\{d \in C \mid aRd\}) \\ &= \sum_{\mathbf{u}_0 \subset \mathbf{u}} \# \{d \in C \mid N_A(d) \supset \{a\}, \mathbf{u}(d) = \mathbf{u}_0\} = k_a \cdot c_1. \end{aligned}$$

Here c_1 is a constant appearing in claim a), and

$k_a = \#\{\mathbf{u}_0 \subset \mathbf{u} \mid (\{a\}, \mathbf{u}_0) \text{ is realisable}\}$. But for $a, b \in A$ $k_a = k_b =: k$: pick i such that $a^{p_i} = b$, then $\{\mathbf{u}_0 \subset \mathbf{u} \mid (\{a\}, \mathbf{u}_0) \text{ is realisable}\}^{x_i} = \{\mathbf{u}_0 \subset \mathbf{u} \mid (\{b\}, \mathbf{u}_0) \text{ is realisable}\}$. (This argument is only needed in the case $m+1 = 2$.) Thus we define the constant $d_{r+1} := k \cdot c_1$.

A is $\mathbf{u}'_c\text{-}\mathcal{K}_m$ -free. Otherwise there would exist $a_1, \dots, a_m \in A$ and $(V_1, \dots, V_r, U_d^{r+1}) \in \mathbf{u}'_c$ such that $a_k R a_l$ (for $1 \leq k < l \leq m$) and $a_k \in V_j$ (for $1 \leq k \leq m$, $1 \leq j \leq r$) and $a_k \in U_d^{r+1}$. But then $(V_1, \dots, V_r) \in \mathbf{u}_c$, $d \in V_j$, $a_k R d$. This means $a_1, \dots, a_m, d$ contradicts the $\mathbf{u}_c\text{-}\mathcal{K}_{m+1}$ -freeness of C .

For $1 \leq i \leq n$ p_i is a χ'_i -permorphism. Furthermore the conditions on the colours $\mathbf{u}_a^{r+1}$ (for $a \in A$) are satisfied, e.g. for $a \in D_i$ we have $(U_a^{r+1})^{x_i} = U_{a^{h_i}}^{r+1} = U_{a^{p_i}}^{r+1}$ because $p_i \subset h_i$.

By the Lemma for m we find a finite $\mathbf{u}'$ -graph B , $B \supset A$, B $\mathbf{u}'_c\text{-}\mathcal{K}_m$ -free and $f_1, \dots, f_n \in \text{Aut}(B)$, $f_i \supset p_i$, f_i a χ'_i -permorphism having the indicated properties. Now we consider B just as a $\mathbf{u}$ -graph. The only thing we still have to check, is that B is $\mathbf{u}_c\text{-}\mathcal{K}_{m+1}$ -free.

Suppose there exist $(V_1, \dots, V_r) \in \mathbf{u}_c$ and elements $a_0, \dots, a_m \in B$ such that $a_k R a_l$ and $a_k \in V_j$. W.l.o.g we suppose $a := a_0 \in A$: Otherwise choose $f \in \langle f_1, \dots, f_n \rangle$, such that $a_0^f \in A$ and choose $\chi' \in \text{Sym}(\mathcal{L}')$ such that f is a χ' -permorphism and let $\chi = \chi' \upharpoonright_{\mathcal{L}}$; now still $(V_1^\chi, \dots, V_r^\chi) \in \mathbf{u}_c$ and $a_k^f R a_l^f$ and $a_k^f \in V_j^\chi$. Because $a \in V_j$ we have $(V_1, \dots, V_r, U_a^{r+1}) \in \mathbf{u}'_c$. The additional condition on B implies now: $a_k \in \mathbf{u}_a^{r+1}$ ($1 \leq k \leq m$). Thus we get a contradiction to the $\mathbf{u}'_c\text{-}\mathcal{K}_m$ -freeness of B . ■(Lemma, Theorem)

Theorem 3 *Let $m \geq 1$, let M be the generic countable $\mathcal{K}_m$ -free graph. M has the Small Index Property*

Proof: The proof is given in [HHLS]. The use there of Hrushovski's Lemma in the proof of the Small Index Property for the generic graph must be replaced by theorem 1. ■(Theorem)

Now we are turning to the case of the digraphs:

Lets suppose there is a fixed (possibly infinite) family $\mathcal{F}$ of finite tournaments (i.e. of digraphs F such that for every $a, b \in F$ if $a \neq b$ then aRb or bRa). Lets look at the class $\mathcal{K}$ of all finite $\mathcal{F}$ -free digraphs A , i.e. of digraphs such that no $F \in \mathcal{F}$ is embeddable into A . This class has the (free) Amalgamation Property, and the resulting generic countable digraph $M_{\mathcal{F}}$ will be called a Henson digraph. Henson [Hen] showed that there continuum many non isomorphic such digraphs.

Theorem 4 a) Let $A \in \mathcal{K}$, let $p_1, \dots, p_n$ be partial isomorphisms on A , then there exist $B \in \mathcal{K}$, $B \supset A$ and $f_1, \dots, f_n$ automorphisms of B , $f_i \supset p_i$.
b) The Small Index Property holds for $M_{\mathcal{F}}$. I.e. it holds for all Henson digraphs.

Proof:

Again The proof given in [HHLS] shows that b) follows from a).

It suffices to prove a) in the case that $\mathcal{F}$ is a finite class: Let $\mathcal{F}$ be infinite and let $A \in \mathcal{K}$. Let $m := \#A$. Let $\mathcal{F}_1$ be a finite family of tournaments of size $m + 1$ containing every isomorphism type of such a tournament. We define $\mathcal{F}_0 := \{F \in \mathcal{F} \mid \#F \leq m\} \cup \mathcal{F}_1$ and we assume w.l.o.g $\mathcal{F}_0$ to be finite. Now A is $\mathcal{F}_0$ -free and every $\mathcal{F}_0$ -free graph B is also $\mathcal{F}$ -free, i.e. in $\mathcal{K}$.

Now we want to prove a) in the case $\mathcal{F}$ is finite by induction on the maximal size of a tournament in $\mathcal{F}$. This prove will be very similar to the prove of theorem 1. We only want to point out the differences. Again the following lemma will be the “permorphism version” of the Theorem, which is provable by induction. Here we will have no restriction on the cardinality of colours a single point can have.

Lemma 5 Let $\mathcal{U}$ be a finite set of colours, let $\chi_1, \dots, \chi_n \in \text{Sym}(\mathcal{U})$. Let $T_1, \dots, T_m$ be finite tournaments, $T_j = \{t_1^j, \dots, t_{l_j}^j\}$, let for $1 \leq j \leq m$ $\mathcal{U}_j \subset (\text{Pot}(\mathcal{U}))^{l_j}$, $\mathcal{U}_j$ χ_i -invariant. Let A be a $\mathcal{U}$ -coloured digraph such that for every j ($1 \leq j \leq m$) A is $\mathcal{U}_j$ - T_j -free (i.e. there does not exist an embedding of digraphs $i : T_j \rightarrow A$, $s_k := i(t_k^j)$ such that $(\mathcal{U}(s_1), \dots, \mathcal{U}(s_{l_j})) \in \mathcal{U}_j$). Let $p_1, \dots, p_n$ be partial mappings such that p_i is a χ_i -permorphism.

Then there exists a finite $\mathcal{U}$ -coloured digraph B , $B \supset A$, B $\mathcal{U}_j$ - T_j -free ($1 \leq j \leq m$) and $f_1, \dots, f_n$ total permorphisms on B , $f_i \supset p_i$.

B satisfies in addition: For every mapping $U : A \rightarrow \mathbf{U}$ (we will write U_a instead of $U(a)$)

if for $b \in A : (bRa \iff b \in U_a)$ (rsp.: $aRb \iff b \in U_a$)

and if for $a \in D_i = D(p_i) : (U_a)^{x_i} = U_{a^{p_i}}$

then for every $b \in B$, for every $a \in A : (bRa \Rightarrow b \in U_a)$ (rsp.: $aRb \Rightarrow b \in U_a$).

Proof:

The proof of the lemma in this version of the preprint will be rather sketchy. The proof goes by induction on the maximal size of the tournaments T_j ($1 \leq j \leq m$) involved. In the case that the maximal size is 1 it goes exactly like the proof in the case of the graphs (case $m = 1$). So let's suppose that the maximal size is > 1 .

Let us first introduce some notation. If A is a digraph and a is a point (possibly element of a digraph extending A) we denote by $N_A^+(a) := \{b \in A \mid aRb\}$ and by $N_A^-(a) := \{b \in A \mid bRa\}$.

Let $T_1, \dots, T_r$ be the tournaments of size > 1 and $T_{r+1}, \dots, T_m$ be the tournaments of size 1. We write for $1 \leq j \leq r$ $T_j^+ := N_{T_j}^+(t_1^j)$ and $T_j^- := N_{T_j}^-(t_1^j)$, thus $T_j - \{t_1^j\} = T_j^+ \cup T_j^-$.

We will call C (where $C \supset A$) free of critical copies of T_j if there is no embedding $\rho : T_j \rightarrow C$ such that (if one denotes $s_l = \rho(t_l^j)$) $s_2, \dots, s_{l_j} \in A$ and such that $(\mathbf{U}(s_1), \dots, \mathbf{U}(s_{l_j})) \in \mathbf{U}_j$.

Now the first step will be, to find $C \supset A$, such that C is free of critical copies of T_j and such that there are bijections $h_1, \dots, h_n \in \text{Sym}(C)$ such that for every $V \in \mathbf{U}$, for every $c \in C : c \in V \iff c^{h_i} \in V^{x_i}$ and such that for every $a \in D_i$, $c \in C : (aRc \iff a^{p_i} R c^{h_i})$ and $(cRa \iff c^{h_i} R a^{p_i})$.

In this case a subset $A_0^+ \subset A$ and a disjoint subset $A_0^- \subset A$ and a set of colours $\mathbf{U}_0$ determine a type over A . We call $(A_0^+, A_0^-, \mathbf{U}_0)$ realisable, if there does not exist $j \leq r$ and an embedding (as digraphs) $i : T_j^+ \cup T_j^- \rightarrow A_0^+ \cup A_0^-$ such that $i(T_j^+) \subset A_0^+$ and $i(T_j^-) \subset A_0^-$ and such that $(\mathbf{U}_0, \mathbf{U}(s_2), \dots, \mathbf{U}(s_{l_j})) \in \mathbf{U}_j$ (again $s_l = i(t_l^j)$). Note that for $C \supset A$: C is free of critical copies of T_j (for every $j \leq r$) iff for every $c \in C : (N_A^+(c), N_A^-(c), \mathbf{U}(c))$ is realizable. Now again choose constants c_t ($1 \leq t \leq \#A$) and choose C to ensure that for every $A_0^+, A_0^- \subset A$, $\mathbf{U}_0 \subset \mathbf{U}$:

$$\#(\{c \in C \mid N_A^+(c) \supset A_0^+, N_A^-(c) \supset A_0^-, \mathbf{U}(c) = \mathbf{U}_0\})$$

$$= \begin{cases} c_{\#(A_0^+ \cup A_0^-)} & \text{if } (A_0^+, A_0^-, \mathbf{u}_0) \text{ is realisable} \\ 0 & \text{otherwise} \end{cases}$$

As in the proof of Lemma 2 one proves that bijections $h_1, \dots, h_n$ with the desired properties exist.

Now our new set of colours will be $\mathbf{u}' = \mathbf{u} \dot{\cup} \{U_c^+ \mid c \in C\} \dot{\cup} \{U_c^- \mid c \in C\}$ (with new colours U_c^+, U_c^-). For $a \in A$ we define ($a \in U_c^+$ iff cRa) and ($a \in U_c^-$ iff aRc). This way we equip A with a $\mathbf{u}'$ -structure and p_i is a χ'_i -permorphism, if we define $(U_c^+)^{\chi'_i} = U_{c^{h_i}}^+$ and $(U_c^-)^{\chi'_i} = U_{c^{h_i}}^-$ (for $1 \leq i \leq n$). For $1 \leq j \leq r$ we set $T'_j = \{t_2^j, \dots, t_{l_j}^j\}$ and we say (for $\mathbf{v}_2, \dots, \mathbf{v}_{l_j} \subset \mathbf{u}'$): $(\mathbf{v}_2, \dots, \mathbf{v}_{l_j}) \in \mathbf{u}'_j$ iff there exists $c \in C$ such that for $2 \leq l \leq l_j$ $U_c^{\epsilon_l} \in \mathbf{v}_l$ and such that $(\mathbf{u}(c), \mathbf{v}_2 \cap \mathbf{u}, \dots, \mathbf{v}_{l_j} \cap \mathbf{u}) \in \mathbf{u}_j$, where $U_c^{\epsilon_l} = U_c^+$, if $t_l^j \in T_j^+$ and $U_c^{\epsilon_l} = U_c^-$, if $t_l^j \in T_j^-$.
 A is $\mathbf{u}'_j$ - T'_j -free.

By induction there is an $\mathbf{u}'_j$ - T'_j -free (for $1 \leq j \leq r$) digraph B , which is also $\mathbf{u}_j$ - T_j -free (for $r < j \leq m$), which satisfies all we want. The only thing which still needs checking is that B is $\mathbf{u}_j$ - T_j -free (for $1 \leq j \leq r$).

Lets suppose there exists a j $1 \leq j \leq r$ and there exists an embedding $i : T_j \rightarrow B$, such that $(\mathbf{u}(s_1), \dots, \mathbf{u}(s_{l_j})) \in \mathbf{u}_j$ (if we write $s_l := t_l^j$). We can suppose that $s_1 \in A$. We consider (for $\epsilon = +$ and for $\epsilon = -$) the mapping $U^\epsilon : A \rightarrow \mathbf{u}'$ such that $U^\epsilon(a) = U_a^\epsilon$, and we fix $c = s_1 \in A$. Take l , $2 \leq l \leq l_j$; if $t_l^j \in T_j^+$, then $t_1^j R t_l^j$, so $c R s_l$, so by the additional property B satisfies: $s_l \in U_c^+$. The same way one checks that if $t_l^j \in T_j^-$, then $s_l \in U_c^-$. Now we consider the embedding i restricted to $T'_j = T_j - \{t_1^j\}$. Now it is easy to check that $(\mathbf{u}'(s_2), \dots, \mathbf{u}'(s_{l_j})) \in \mathbf{u}'_j$. But this is in contradiction to the $\mathbf{u}'_j$ - T'_j -freeness of B . ■(Lemma, Theorem)

Finally we want to mention that there is a more general theorem stating the possibility of extending partial isomorphisms. It is a theorem about relational structures in any relational language. To state it we need some notation. Let S be a finite relational language.

Definition:

- Let L be a S -structure. L is called a link structure if it consists of just one element or if there is a k -ary ($k \in \omega$) relation symbol R in S and a k -tuple $(a_1, \dots, a_k)$ in L such that $Ra_1 \dots a_k$ and such that $a_1, \dots, a_k$ contains all elements of L .
- Let A and B be S -structures. A and B have the same link type if for every link structure L : L is embeddable into A iff L is embeddable into B .
- Let T, A be S -structures, let $\rho : T \rightarrow A$ be a function. ρ is called a weak homomorphism (notation: $\rho : T \rightarrow_w A$) if for every R in S (R k -ary) and every $s_1, \dots, s_k \in T$: If $Rs_1 \dots s_k$ (in T) then $Rs_1^\rho \dots s_k^\rho$ (in A).
- Let $\mathcal{F}$ be a set of finite S -structures. Let A be a S -structure. A is called $\mathcal{F}$ -free if there does not exist $T \in \mathcal{F}$ and $\rho : T \rightarrow_w A$.

For example if S consists of just one binary relation symbol, and A is a graph (rsp. digraph) and if the S -structure B has the same link structure as A then B is a graph (rsp. digraph); but this is not true for tournaments.

Theorem 6 *Let $\mathcal{F}$ be a finite set of finite S -structures. Let A be a finite $\mathcal{F}$ -free S -structure. Let $p_1, \dots, p_n$ be partial isomorphisms on A . There exist a finite $\mathcal{F}$ -free S -structure B , $A \subset B$ and automorphisms $f_1, \dots, f_n$ on B ($f_i \supset p_i$) such that B and A have the same link type.*

The proof of the theorem is just the translation of the proofs of theorem 1 and theorem 4 a) into this more general context. Note that a weak homomorphisms mapping $\mathfrak{K}_m$ to a graph A (and a weak homomorphisms mapping a tournament into a digraph) is necessarily an embedding.

References

- [C] P. Cameron, Oligomorphic Permutation Groups, LMSLNS 152,
Cambridge University Press, 1990,
- [Hg] B.Herwig, Extending Partial Isomorphisms, Combinatorica, to appear,
- [Hr] E. Hrushovski, Extending Partial Isomorphisms of Graphs,
Combinatorica 12(1992), 411-416
- [HHLS] Hodges, Hodkinson, Lascar, Shelah: The Small Index Property for
 ω -stable, ω -categorical, structures and for the random graph,
Journal of the LMS 48(1993), 204-218.
- [T] J.K. Truss, Generic Automorphisms of Homogeneous Structures,
preprint, 1990.